

Q&A 104.4

Can small volume intramuscular injections be given to patients taking oral anticoagulants?

Prepared by UK Medicines Information (UKMi) pharmacists for NHS healthcare professionals
Before using this Q&A, read the disclaimer at www.ukmi.nhs.uk/activities/medicinesQAs/default.asp
Date prepared: 5th October 2015

Background

As increasing numbers of people are prescribed oral anticoagulants, for conditions such as atrial fibrillation, it is likely that a number will require a small volume intramuscular (IM) injection at some point. Factors that need to be considered in these cases include:

- Is there an increased risk of adverse effects, such as bruising or haematoma?
- Could the injection affect anticoagulant control?
- Could the anticoagulant alter the effectiveness of the injection e.g. vaccines?

This Medicines Q&A focuses on the potential risks of adverse effects associated with the administration of small volume IM injections to patients taking oral anticoagulants. Detailed analyses of drug interactions are outside the scope of this Q&A and further information should be sought on an individual patient basis.

Answer

Most of the published studies available on this subject relate to IM influenza vaccination in patients taking older anticoagulants, such as warfarin. Data are limited and some findings are contradictory. Over recent years there has been a wider uptake of the newer oral anticoagulants, apixaban, rivaroxaban, dabigatran and edoxaban tosylate, collectively named novel oral anticoagulants (NOACs), but there is very little published information about the risks of adverse effects with NOACs and IM injections.

Influenza vaccination

People taking oral anticoagulants commonly require vaccination against the influenza virus.

Studies of oral anticoagulants and intramuscular influenza vaccination

- A randomised, single-blinded trial (n=229) compared subcutaneous (SC) to IM influenza vaccine administered in the deltoid area in patients on oral anticoagulants (98% acenocoumarol, 2% warfarin) (1). Skin lesions (such as erythema) 24 hours after injection were reported by 37.4% of the SC group and 17.4% of the IM group (number needed to harm 5 [95% confidence interval 3.1 to 11.3]). No major side effects were reported. Two patients in the SC group and one in the IM group experienced a haematoma. No statistically significant differences were seen in international normalised ratio (INR) results between the groups.
- A randomised, double-blind, cross-over trial included 104 patients on stable warfarin therapy (2). Patients either received influenza vaccine or placebo into the deltoid muscle at the start of the first 28 day period and then, after a 14-day washout, crossed over to the alternative treatment and were followed-up for a further 28 days. Mean INR results, weekly warfarin dose and time in the therapeutic range were similar between the two groups in the two study periods. No major bleeding events were seen. Antibody response was not affected by warfarin treatment. The authors concluded that patients on stable doses of warfarin can safely be given influenza vaccine without the need to increase the frequency of INR testing.

- A study of 41 male patients taking warfarin assessed the effects of a 0.5mL influenza vaccine as a single IM injection in the deltoid region, followed by application of firm pressure for 5 minutes (3). All patients were followed up for 14 days and there were no cases of localised bleeding or any change in arm girth after injection. Vaccination did not have a significant effect on prothrombin time.
- Thirteen male patients received 0.5mL influenza vaccine intramuscularly and 13 received the same vaccine subcutaneously in the deltoid region in a single-blind study (4). Three patients in each group had discomfort or pain at the injection site, but no local bleeding was recorded. Ten recipients of IM injection were followed up by phone 2 days later and no bleeding, bruising or swelling was described. The route of administration did not affect serum influenza antibody titres.
- A case-control study during the 2001-02 influenza season looked at the effect of influenza vaccination on prothrombin time in patients receiving long-term oral anticoagulant treatment (98% warfarin, 2% acenocoumarol) (5). Ninety patients [mean INR 2.79 (\pm 0.83)] who received IM influenza vaccine were compared to 45 matched controls who did not receive vaccination. Influenza vaccination was associated with an average increase in INR of 0.56 seven to ten days after vaccination. Two patients experienced bleeding episodes (epistaxis and muscular haematoma). The authors suggest that INR should be monitored carefully in anticoagulated patients in the period immediately following influenza vaccination.
- A systematic review published in 2012 attempted to assess the validity of the assertion that the influenza vaccine may potentially interfere with the anticoagulant efficacy of long-term warfarin therapy (6). The review did not indicate any consistent, clinically relevant effect of influenza vaccine on INR results. The review also discussed that there does not appear to be an increased risk of haematoma following IM administration of the influenza vaccines.

Other information sources

- A resource providing information and advice about drug interactions discusses that in small to medium studies (some uncontrolled), most did not demonstrate a significant change in prothrombin time or INR in patients taking **coumarins**, such as **warfarin** (7). Some studies found slight increases or decreases in anticoagulant effect, but these are probably of limited clinical relevance. However, in one large observational case-control study, an increase in INR from 2.64 to 3.85 was seen in about half of the 90 patients and 2 had bleeding episodes. The reasons for the difference between this study and others are not known. Overall, the weight of evidence suggests that the concurrent use of warfarin and influenza vaccination is usually safe and uneventful, but it would be prudent to be alert as there have been some reports of bleeding attributed to an interaction. **Acenocoumarol** does not normally interact with influenza vaccination.
- The British National Formulary (BNF) states that the anticoagulant effect of **warfarin** can possibly be enhanced by **influenza vaccine**, but serious consequences are not expected (8).
- An American reference source (AHFS Drug Information) also states that **influenza vaccine** may increase a patient's response to **coumarin-type anticoagulants** (9).

Practical Advice

Most seasonal influenza vaccines are licensed for both IM and deep subcutaneous (SC) use. An NHS Clinical Knowledge Summary recommends that **seasonal influenza vaccine** is given by deep SC injection if patients are taking warfarin (10). It has been suggested that giving **influenza vaccine** to patients taking **warfarin** is normally safe (7). However, due to the theoretical risk of local muscle haematoma, it may be preferable to give such injections by deep SC injection to patients taking **coumarins** and **related anticoagulants**. There is no advice on this subject in the manufacturer's Summary of Product Characteristics (SPC) for the original brand of warfarin (Marevan®) (11).

Other vaccines

Limited information on the administration of vaccines to patients receiving oral anticoagulants is available. Individual SPCs should be consulted for further information. For example, the manufacturers of **yellow fever vaccine** (Stamari[®]) recommend that due to the risk of injection site haematoma with IM injection, the vaccine should be administered via the SC route for patients taking an anticoagulant (12).

The results of influenza vaccination studies cannot be generalised to other vaccines or to IM injections with a higher volume or with needles of a greater calibre (1,13). The risk-benefit ratio for persons at risk of adverse effects, such as bruising or haematoma, following IM injection must be evaluated by healthcare professionals.

Results from a retrospective cohort study in patients stabilised on **warfarin** suggest that no alteration in anticoagulant control would be expected after use of vaccines such as **23-valent pneumococcal polysaccharide vaccine** or **hepatitis A vaccine** (14).

The safety and efficacy of the combined **tetanus-diphtheria vaccine** administered intramuscularly to anticoagulated patients is the subject of an ongoing Spanish study which is due to publish results in 2016. (15)

No data have been identified that show changes to antibody titres to other vaccines due to oral anticoagulants.

Novel oral anticoagulants

As discussed above, data on the safety of IM drug administration to patients taking NOACs are very limited.

A study in Germany evaluated peri-interventional management of NOACs in unselected patients from daily care (16). Safety data was collected from a prospective registry of over two thousand patients taking a NOAC who also underwent interventional procedures that were classified as either 'minimal', 'minor' or 'major', based on severity of tissue trauma and the risk of peri-procedural bleeding. Intramuscular injections were classified as **minor** procedures which involved little tissue trauma but a 'relevant bleeding risk'. To put this risk classification into perspective; procedures such as cataract surgery and pace-maker related surgeries were also deemed as 'minor' interventions. The authors concluded that due to the pharmacological profile of the NOACs (short half-life and fast onset of action) continuation or short-term interruption of a NOAC, without use of bridging therapy, is a safe strategy for most patients undergoing **non –major** invasive procedures, but advised that it would be prudent to conduct a careful risk: benefit evaluation before any patient specific decisions are made (16)

Expert opinion from a consultant haematologist supports this advice, suggesting that in clinical practice, if time allows, it is desirable to administer an IM injection 24 hours after previous NOAC dose and then to restart the NOAC the following day, assuming that there are no signs of haematoma or bruising. Although, in the opinion of the haematologist, short term interruption is unlikely to pose a high risk of a thrombotic event occurring, each patient's individual circumstances needs to be carefully evaluated on a case-by-case basis (17).

The Summary of Product Characteristics (SPC) for **dabigatran etexilate** states that injection site haemorrhage occurs rarely (18). Vessel puncture site haematoma is seen uncommonly in patients taking **apixaban** (19). Similar data are not included in the SPC for **rivaroxaban**. (20) and edoxaban (21)

Other intramuscular drugs

Comprehensive information about the administration of individual IM drugs to patients receiving oral anticoagulants is beyond the scope of this Medicines Q&A. Individual SPCs should be consulted for further information. For example, some SPCs for IM drugs state that this route has not been evaluated in patients receiving anticoagulants because of the risk of haematoma. Therefore, the risk-benefit ratio for persons at risk of adverse effects, such as bruising or haematoma, following IM injection must be evaluated by healthcare professionals.

Actions when considering a small volume IM injection for a patient taking an oral anticoagulant:

- ◆ If possible, avoid IM injections, even when the INR is being closely controlled, and do not use if the INR is raised above the therapeutic range;
- ◆ If small volume IM injections are necessary:
 - check the relevant SPC;
 - evaluate the risk-benefit ratio for each patient;
 - administer injections into an upper extremity as a precaution to permit easy access for manual compression, inspection of bleeding, and/or application of pressure bandages if necessary (9);
 - use a fine needle and apply firm pressure for at least 2 minutes immediately after vaccination (13)
 - Inform the patient that they should watch out for localised bleeding.

Summary

- ◆ Limited information is available on the administration of small volume intramuscular (IM) injections to patients taking oral anticoagulants, especially the newer Novel Oral Anticoagulants (NOACs). Most published studies relate to older anticoagulants, such as warfarin.
- ◆ IM injections should be avoided where possible, even when the INR is being closely controlled, and not used if the INR is raised above the therapeutic level.
- ◆ Healthcare professionals should weigh up the benefits and potential risks on an individual patient basis.
- ◆ An IM site on an upper extremity should be preferred to a central site, so that any bleeding can be seen, and compression applied if necessary.
- ◆ Most seasonal influenza vaccines are licensed for both IM and deep SC use. It is often recommended that **seasonal influenza vaccine** is given by deep SC injection if patients are taking **warfarin** or related anticoagulants).
- ◆ Studies of influenza vaccination in patients taking oral anticoagulants have shown conflicting results. Some note that patients on stable doses of warfarin do not need to increase the frequency of INR testing, whereas some patients have experienced bleeding or haematoma.
- ◆ Currently, data on NOACs are very limited, although the manufacturers of some NOACs warn of the risk of injection site haematoma or haemorrhage.
- ◆ A single study from Germany suggests that continuation or short-term interruption of a NOAC, without use of bridging therapy, is possibly a safe strategy for most patients undergoing **non – major** invasive procedures, such as intramuscular injections. Conducting a careful risk: benefit evaluation before any patient specific decisions are made is still a prudent course of action.

Limitations

- The information above is based upon limited published data
- The list of drugs is not comprehensive. Drugs which have not been included cannot be assumed to be safe when administered to patients taking oral anticoagulants.
- Detailed analyses of drug interactions are outside the scope of this Q&A and further information should be sought on an individual patient basis.
- Only the use of small volume IM injections has been considered. The use of other IM injections e.g. depot injections is beyond the scope of this Medicines Q&A.

References

1. Casajuana J, Iglesias B, Fàbregas M et al. Safety of intramuscular influenza vaccine in patients receiving oral anticoagulation therapy: a single blinded multi-centre randomized controlled clinical trial. *BMC Blood Disorders* 2008;8:1.
2. Iorio A, Basileo M, Marcucci M et al. Influenza vaccination and vitamin K antagonist treatment. A placebo-controlled, randomized, double-blind crossover study. *Arch Intern Med* 2010;170:609-16.
3. Raj G, Kumar R, McKinney WP. Safety of intramuscular influenza immunization among patients receiving long-term warfarin anticoagulation therapy. *Arch Intern Med* 1995;155:1529-31.
4. Delafuente JC, Davis JA, Meuleman JR et al. Influenza vaccination and warfarin anticoagulation: a comparison of subcutaneous and intramuscular routes of administration in elderly men. *Pharmacotherapy* 1998;18:631-6.
5. Paliani U, Filippucci E, Gresele P. Significant potentiation of anticoagulation by flu-vaccine during the season 2001-2002. Letter. *Haematol* 2003;88:599-600.
6. Kuo, A.M, Brown, J.N, Clinard, V. Effect of influenza vaccination on international normalized ratio during chronic warfarin therapy. *Journal clinical pharmacy and therapeutics*. 2012, 37, 505-509
7. Coumarins and related drugs + vaccines; influenza. Baxter K, editor. *Stockley's Drug Interactions: A source book of interactions, their mechanisms, clinical importance and management*. Electronic edition. London: Pharmaceutical Press. Accessed via <http://www.medicinescomplete.com/> on 2nd October 2015
8. Joint Formulary Committee. *British National Formulary*. Electronic edition. London: BMJ Group and Pharmaceutical Press. Accessed via www.medicinescomplete.com on 2nd October 2015
9. Warfarin Sodium monograph. AHFS Drug Information. USA (Bethesda): American Society of Health-System Pharmacists, Inc. Accessed via www.medicinescomplete.com on 2nd October 2015
10. Clinical Knowledge Summary (CKS). National Institute for Health and Care Excellence (NICE). Immunizations - seasonal influenza – Management. Accessed from <http://cks.nice.org.uk/immunizations-seasonal-influenza> on 2nd October 2015.
11. Summary of Product Characteristics – Marevan 5mg tablets (warfarin sodium). Amdipharm Mercury Company Ltd. Date last updated on eMC: 26th May 2014. Accessed via www.medicines.org.uk on 2nd October 2015
12. Summary of Product Characteristics – Stamaril (yellow fever vaccine). Sanofi Pasteur MSD Ltd. Date last updated on eMC 18th September 2013. Accessed via www.medicines.org.uk on 2nd October 2015
13. Ringwald J, Strobel J, Eckstein R. Travel and oral anticoagulation. *Journal of Travel Medicine* 2009;16:276-83.
14. Jackson ML, Nelson JC, Chen RT et al for the Vaccine Safety Datalink investigators. Vaccines and changes in coagulation parameters in adults on chronic warfarin therapy: a cohort study. *Pharmacoepidemiol Drug Safety* 2007;16:790–6.
15. Lago-Deibe Fernando I, Martin- Miguel, M-V. et al. The safety and efficiency of the tetanus vaccine intramuscularly versus subcutaneously in anticoagulated patients: a randomized clinical trial. Study Protocol. Trial registration ISRCTN69942081. *BMC Family Practice* 2014; 15:147.
16. Beyer–Westendorf J, Gelbricht V, Forster K, Ebertz F, et al. Peri-interventional management of novel oral anticoagulants in daily care: results from the prospective Dresden NOAC registry. *European Heart Journal* 2014;35(28):1888-96. Full text available at <http://eurheartj.oxfordjournals.org/content/ehj/35/28/1888.full.pdf>
17. Author's personal communications with a consultant haematologist, Clinical lead, Southern Haemophilia Network. – September 2015
18. Summary of Product Characteristics – Pradaxa 150mg hard capsules (dabigatran etexilate mesilate). Boehringer Ingelheim Ltd. Date last updated on eMC 2nd September 2015. Accessed via www.medicines.org.uk on 2nd October 2015.
19. Summary of Product Characteristics – Eliquis 2.5mg tablets(apixaban). Bristol-Myers Squibb-Pfizer. Date last updated on eMC: 1st October 2015. Accessed via www.medicines.org.uk on 2nd October 2015.

20. Summary of Product Characteristics – Xarelto 10mg Film-Coated tablets (rivaroxaban). Bayer plc. Date last updated on eMC: 1st July 2015. Accessed via www.medicines.org.uk on 2nd October 2015.
21. Summary of Product Characteristics – Lixiana 15mg Film-Coated tablets (edoxaban tosilate). Daiichi Sankyo UK Ltd. Date last updated on eMC: 2nd July 2015. Accessed via www.medicines.org.uk on 8th October 2015.

Quality Assurance

Prepared by

Stephen Fleck (based on earlier work by Simon Wills and Sue Gough), Lead Medicines Advice Pharmacist, Southampton Medicines Advice Service, University Hospital Southampton NHS Foundation Trust.

Date Prepared

5th October 2015

Checked by

Samantha Owen, Principal Pharmacist Critical Evaluation, Southampton Medicines Advice Service, University Hospital Southampton NHS Foundation Trust.

Date of check

1st December 2015

Search strategy

- Medline
 - exp WARFARIN/ and exp INJECTIONS, INTRAMUSCULAR/
 - exp ACENOCOUMAROL/ and exp INJECTIONS, INTRAMUSCULAR/
 - exp PHENINDIONE/ and exp INJECTIONS, INTRAMUSCULAR/
 - exp ANTITHROMBINS/ and exp INJECTIONS, INTRAMUSCULAR/
 - dabigatran.ti,ab and exp INJECTIONS, INTRAMUSCULAR/
 - apixaban.ti,ab and exp INJECTIONS, INTRAMUSCULAR/;
 - rivaroxaban.ti,ab and exp INJECTIONS, INTRAMUSCULAR/
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp WARFARIN/
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp ACENOCOUMAROL/
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp PHENINDIONE/
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp ANTITHROMBINS/
 - exp VACCINATION/ or exp IMMUNIZATION/ and dabigatran.ti,ab
 - exp VACCINATION/ or exp IMMUNIZATION/ and rivaroxaban.ti,ab
 - exp VACCINATION/ or exp IMMUNIZATION/ and apixaban.ti,ab
 - exp VACCINATION/ or exp IMMUNIZATION/ and edoxaban.ti,ab
- EMBASE
 - exp WARFARIN/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - [Limit to: Human and English Language]
 - exp ACENOCOUMAROL/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - exp PHENINDIONE/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - exp ANTITHROMBIN/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - exp DABIGATRAN ETEXILATE/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - exp APIXABAN/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - exp RIVAROXABAN/ and exp INTRAMUSCULAR DRUG ADMINISTRATION/
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp WARFARIN/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp ACENOCOUMAROL/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp PHENINDIONE/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp ANTITHROMBIN/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp DABIGATRAN ETEXILATE/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp RIVAROXABAN/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp APIXABAN/ (limited to Human)
 - exp VACCINATION/ or exp IMMUNIZATION/ and exp EDOXABAN/ (limited to Human)
- In-house databases and resources